



## Foliar Pathogens in Guam: *Fusarium* (*Gibberella*)

Diseases: Fusarium Wilt, Leaf Spot of Orchid

Robert L. Schlub, Ph.D., Julia Hudson, and Elizabeth Hahn  
Cooperative Extension & Outreach, College of Natural & Applied Sciences,  
University of Guam




Figure 1. [A] Spores of *Fusarium* produced by short conidiophores.  
[B] Macroconidia

Source: (Streets, R. 1972). *The Diagnosis of Plant Diseases: A Field and Laboratory Manual Emphasizing the Most Practical Methods for Rapid Identification.*

### Introduction

Most species of *Fusarium* are widely distributed, soilborne, non-pathogenic, and active as soil decomposers; however, it is pathogenic on several hundred plant species including economically important food crops such as sweet potatoes, tomatoes, legumes, melons, and bananas (in which the infection is known as Panama disease). As a pathogen, it is most frequently reported to cause root and stem rot and vascular wilt diseases. Less frequently are reports of fruit rot and foliar diseases. The majority of the *Fusarium* wilt diseases are caused by *Fusarium oxysporum*. Fortunately, it exists as host-specific biotypes; therefore, wilt of watermelon on Guam will not spread to cucumbers, tomatoes, or any other crops. *Fusarium proliferatum* is a major pathogen of orchids on Guam. It causes black and yellow leaf spots on *Cymbidium*, *Ondontioda*, *Dendrobium*, and *Cattleya*; and stem and root rots on *Dendrobium*, *Cymbidium* and *Phalaenopsis*.

### Hosts

In the Index of Plant Diseases on Guam *Fusarium* was mentioned on 21 hosts, causing foliar symptoms in 13 of those hosts. This includes some of Guam's major agronomic crops such as watermelon, cucumbers, melon, banana, tomatoes, peppers, and cabbage. In the Diseases of Cultivated Crops in Pacific Island Countries. It was listed as a foliar pathogen on banana, coffee, gerbera, sweet pepper, and peach.

### Morphology of *Fusarium* spp.

The genus is large and taxonomically difficult because it is notoriously variable and ubiquitous. Three types of asexual spores are produced by most *Fusarium* spp.: macroconidia (Fig. 3A), microconidia (Fig. 3B), and chlamyospores (Fig. 3 C,D). Mycelium of most *F. oxysporum* species is white and sparse under normal conditions. Short hyaline conidiophores are produced in a sporodochium (cluster of conidiophores) which grows on the surface of infected tissue, which may appear as a salmon-pink crust (Fig. 2). Conidiophores are 10-20  $\mu\text{m}$  and branching with conidia produced singularly at the tips (Fig. 1A). Macroconidia are commonly found. Macroconidia can vary greatly in shape and size, but generally are 3-5 x 15-44  $\mu\text{m}$ , colorless, multicellular (3-5 cells), curved ("sickle" or "canoe" shaped), have 3-5 septa (cross-walls), with an apical cell and foot-shaped basal cell (a constricted area where the spore was attached during formation) (Fig. 3A). Microconidia are colorless, rounded, 1-2 celled, and small (3-4 x 5.5 12  $\mu\text{m}$ ) (Fig. 3B). *Fusarium proliferatum* is characterized by producing microconidia in chains. Chlamyospores are thick walled with rounded cells typically found in older mycelium, and are the fungus' survival structure (Fig. 3 C,D).




Figure 2. Salmon-pinkish crust of tightly packed spores, typical of *Fusarium*  
Photo: R.L. Schlub






Figure 3. [A] Macroconidia, [B] Microconidia, and [C, D] Chlamydospores of *Fusarium oxysporum*. Scale bars= 10  $\mu$ m  
Source: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7085860/>

### Visibility of *F. oxysporum*

- **With the unaided eye:** vines and leaves of a diseased plant appear wilted (Fig. 4, 5).
- **With a 14X coddington hand lens:** a cottony crust of mycelium and clumps of tightly packed spores (sporodochium) may be visible on plant surfaces (Fig. 2).
- **With a dissecting microscope:** macroconidia may be seen (Fig. 3A).
- **With a compound microscope:** microconidia, macroconidia, and chlamydospores are clearly visible if present (Fig. 3).

### Disease Development

*Fusarium* fungus lives in the soil on crop residue and weed hosts. It can survive in the plant for many years, and any plant part that comes in contact with the soil can be susceptible to infection, which can occur at any stage of growth. Movement of infected soil by wind, rain, and equipment are the main means of spread of *Fusarium* wilt. The disease is most severe in light, sandy soils with a pH of 5.5– 6.5 with low to moderate soil moisture. Optimum temperature is 28-32°C (82.4-89.6°F). Several species of *Fusarium* are associated with orchids; some are pathogenic and cause symptoms of flower spots, leaf or sheath blights, pseudostem or root rots, wilts, and leaf spots (Fig. 6)

### Foliar Symptoms

“Wilt” is the symptom used to describe a plant that is drooping from lack of water without any outward sign of infection. At first, only a few vines will show symptoms or only one side of the plant will show symptoms (unilateral development) (Fig. 4). Plants will show what appears to be normal midday drooping (Fig. 5) but are slow to recover and eventually may die. Plants that are not killed show

yellowing, stunting, and poor fruit development. Symptoms are most severe during droughts and fruit formation. Symptoms of *Fusarium* spp. foliar infections of orchids include leaf spots and leaf blotch. The infection spreads worldwide because of international trade and monoculture. Black, small spots occur on leaves at the initial stage of infection (Fig. 6), followed by the spots enlarging and leaves turning yellow.



Figure 4. Unilateral wilting caused by *Fusarium*, shown on watermelon  
Source: <https://plantpath.ifas.ufl.edu/u-scout/cucurbit/fusarium-wilt---watermelon.html>



Figure 5. *Fusarium* causing yellowing and wilting on cucumber vines  
Source: <https://www.nexles.com/articles/cucumber-cucumis-sativus-treatments-common-diseases-pests-vegetable/>



Figure 6. *Fusarium* causing leaf spot on orchid  
Source: [http://www.extento.hawaii.edu/kbase/crop/Type/f\\_prolif.htm](http://www.extento.hawaii.edu/kbase/crop/Type/f_prolif.htm)





### For further information

Contact the College of Natural & Applied Sciences, Extension and Outreach at 735-2080 for help or more information. Additional publications can be found on our website at: [uog.edu/extension/publications](http://uog.edu/extension/publications).

### Acknowledgments

This material is based upon work that is supported by the University of Guam and the National Institute of Food and Agriculture, U.S. Department of Agriculture, under award number 2019-38640-29880 through the Western Sustainable Agriculture Research and Education program under project number WPDP20-001. USDA is an equal opportunity employer and service provider. Any opinions, findings, conclusions, or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the view of the U.S. Department of Agriculture or the University of Guam. The factsheet was prepared by UOG Extension Specialist Robert L. Schlub, Ph.D. and UOG Extension Associates Julia Hudson and Elizabeth Hahn on 10/19/2022. Reviewed by Drs. Marin T. Brewer and Leilani G. Sumabat.

### References

Schlub, R.L. (2017). Index of Plant Diseases in Guam. Guam Cooperative Extension, College of Agriculture and Life Sciences.

Kohler, F., Pellegrin, F., Jackson, G., & McKenzie, E. (1997). Diseases of Cultivated Crops in Pacific Island Countries. South Pacific Commission.

Yudin, L. and Schlub, R. eds. (1998) Guam cucurbit guide, (2nd ed 2014). University of Guam UOG Station: Mangilao, Guam 20-33 <http://cnas-re.uog.edu/wp-content/uploads/2016/06/Guam-Cucurbit-Guide-2014-2nd-Edition-optimized.pdf>

Center of Agriculture and Biosciences International (2021). *Fusarium oxysporum f.sp. niveum* (Fusarium wilt of watermelon). CABI Invasive Species Compendium. <https://www.cabi.org/isc/datasheet/24668>

Lobard, L., Sandoval-Denis, M., Lamprecht, S.C., Crous, P.W. (2019). Epitypification of *Fusarium oxysporum*—clearing the taxonomic chaos. *Persoonia*. 43:1-47. DOI: 10.3767/persoonia.2019.43.01. PMID: 32214496; PMCID: PMC7085860

Published: 19 October 2022



Published by the College of Natural & Applied Sciences (CNAS), University of Guam, in cooperation with the U.S. Department of Agriculture, under Dr. Lee S. Yudin, Director/Dean. University of Guam, CNAS, UOG Station, Mangilao, Guam 96923. © For reproduction and use permission, contact [cnasteam@triton.uog.edu](mailto:cnasteam@triton.uog.edu), (671) 735-2080. The University of Guam is an equal opportunity/affirmative action institution providing programs and services to the people of Guam without regard to race, sex, gender identity and expression, age, religion, color, national origin, ancestry, disability, marital status, arrest and court record, sexual orientation, or status as a covered veteran. Find CNAS publications at [uog.edu/extension/publications](http://uog.edu/extension/publications). If you anticipate needing any type of reasonable accommodation or have questions about the physical access provided, please call the UOG EEO/ADA/Title IX Office at (671) 735-2971/2244 or email [efgogue@triton.uog.edu](mailto:efgogue@triton.uog.edu).

